

Spiraea tomentosa in earlier years. But the occurrence on East Wolf Island of *Salix pyrifolia* and *Geocaulon lividum* add support to our present feeling that the flora of the Wolves is characterized by a pronounced boreal influence.

Also, we are appending herewith a list of added taxa on Gull Rock, Wolf Islands, which as noted earlier (p. 96) we had not been able to finish because of adverse weather conditions in 1962. These now make Gull Rock as well surveyed as the other islands.

ADDITIONS TO THE VASCULAR FLORA OF GULL ROCK

- Festuca rubra* L. var. *commutata* Gaudin
Puccinellia laurentiana Fern. & Weath. var. *laurentiana*
Puccinellia paupercula (Holm) Fern. & Weath. var. *alaskana* (Scribn. & Merr.) Fern. & Weath.
Agropyron repens (L.) Beauv. f. *trichorrhachis* Rohlena
Lathyrus japonicus Willd. var. *pellitus* Fern.
Aster novi-belgii L.

This brings the total taxa on Gull Rock to 35.

DEPARTMENTS OF BOTANY AND HORTICULTURE, UNIVERSITY
OF NEW HAMPSHIRE.

FASCIATION IN *EMPETRUM NIGRUM* — Fasciation is a commonly observed yet still little understood abnormal growth pattern. It is worth recording its occurrence in a previously unreported species as a matter of record. *Empetrum nigrum* with fasciated growth in typical cockscomb form was collected on Great Wass Island, Beals, Washington County, Maine September 1963 and has been deposited in the Herbarium of the University of New Hampshire.

R. B. PIKE, UNIVERSITY OF NEW HAMPSHIRE